

# MISCELLANEOUS PAPERS

\*\*\*\*\*  
Centre for Entomological Studies Ankara  
\*\*\*\*\*

No 76

24 08 2001

ISSN 1015-8235

## Contributions to the *Orthoptera fauna of the Middle Taurus Mountains* (South Turkey)

Mustafa Ünal<sup>1</sup>

**Abstract:** Contributions to the *Orthoptera* fauna of the Middle Taurus Mountains (South Turkey).  
*Cent. ent. Stud. Misc. Pap.* 76: 1-8.

The Orthoptera material collected in the Middle Taurus Mountains is evaluated. Totally, 57 species and subspecies are listed. Original references, collecting data and distributional information are given to each species. Remarks are added to some species.

**Key Words:** *Orthoptera*, fauna, Taurus Mountains, Turkey.

The studied area is rather poorly known in the orthopterological literature. The material studied has been collected mainly from Bozkır district in Konya province, Alanya district in Antalya province and Geyik Mountain in the Taurus Mountain Range, situated between both districts. Separately, several specimens have been captured near Tuz Gölü in the Middle Anatolia. Field studies were made in the second half of June and the first quarter of September in 1997. Totally, 604 specimens, belonging to 57 species and subspecies, have been captured by the author from different 35 localities. Among them, 19 species are endemics for Turkey. *Isophya major* Br.-W., *Poecilimon davisi* Karabağ, *Parapoecilimon antalyensis* Karabağ, *Drymadusa limbata grandis* Karabağ, *Parapholidoptera antaliae* Nadig and *Paranocarodes aff. fieberi* Br.-W. are found from Taurus Mountains only and its neighbouring. The genus *Parapoecilimon* Karabağ, with its sole species *P.antalyensis* Karabağ and *Glyphotmethis holtzi spinosus* Karabağ are recorded first time after their original descriptions. Some immature specimens are not listed as they could not be determined with certain in the genera *Tettigonia*, *Bucephaloptera*, *Saga* and *Paranocaracris*. All the specimens studied are in the collection of the author.

### LIST OF SPECIES

#### *Tettigoniidae: Phaneropterinae*

*Acrometopa syriaca* Brunner von Wattenwyl, 1878  
*Acrometopa syriaca* Brunner von Wattenwyl, 1878,  
Mon.Phan., p. 85, 87.

Material Studied: Antalya: Alanya, Alarahan, 75m., 21.6.1997, 1 ♂; Mahmutlar, 100m., 22.6.1997, 1 ♂ nymph; 10 km. N. of Mahmutlar, 22.6.1997, 1 ♂, 1 ♀ (nymphs). Distribution: Greece, Anatolia, Middle East (Ramme, 1951; Bei-Bienko, 1954; Harz, 1969).

*Tylopsis lilifolia* (Fabricius, 1793)  
*Locusta lilifolia* Fabricius, 1793, Ent.Syst., 2:36.

Material Studied: Antalya: Alanya, Alarahan, 75m., 21.6.1997, 2 ♂ nymphs; Mahmutlar, 100m., 22.6.1997, 1 ♀ nymph. Distribution: N. Africa, Europe, Balkans, Anatolia, Middle East, Caucasus, W. Asia (Ramme, 1951; Bei-Bienko, 1954; Harz, 1969).

#### *Isophya karabagi* Uvarov, 1940

*Isophya karabagi* Uvarov, 1940, Ann. mag. Nat. Hist., 4:552.

Material Studied: Konya: Kulu, 1000m., 19.6.1997, 2 ♂, 3 ♀. Distribution: Middle Anatolia (Bei-Bienko, 1954; Karabağ, 1958).

Remarks: This species was known from North of the Middle Anatolia in Ankara province so far. Specimens have been found from West of Tuz Gölü.

<sup>1</sup> Abant İzzet Baysal Üniversitesi, Fen-Edebiyat Fakültesi, Biyoloji Bölümü, 14280 Bolu, Turkey.  
e-mail: [unal@abant.ibu.edu.tr](mailto:unal@abant.ibu.edu.tr)

***Isophya major* Brunner von Wattenwyl, 1878**

*Isophya major* Brunner von Wattenwyl, 1878,  
Mon. Phan., p. 70.

Material Studied: Konya: Bozkır, Kuruçay, 1400m.  
14.6.1997, 4 ♂; 6 km.N.E. of Bozkır, 1400m.  
15.6.1997, 2 ♀, 20.6.1997, 1 ♀; Yalihüyük yaylası,  
1700m. 14.6.1997, 1 ♂; Sorkun, 1200m. 14.6.1997, 2  
♂, 2 ♀ (in alcohol); 3 km.S. of Elmaağac, 1200m.  
13.6.1997, 3 ♂, 2 ♀ (in alcohol); Hadim, 1200-1300m.  
13.6.1997, 2 ♂, 1 ♀; Antalya: Akseki, 1700m.  
24.5.1997, 5 ♂, 7 ♀, 1 ♀ nymph. Distribution: S.  
Anatolia (Ramme, 1951; Bei-Bienko, 1954; Karabağ,  
1958).

***Isophya aff. pentheri* Ramme, 1951**

*Isophya pentheri* Ramme, 1951, Mitt.Zool.Mus.Berlin,  
27: 167.

Material Studied: Konya: Seydişehir, Gökhüyük,  
1000m. 14.6.1997, 2 ♂. Distribution: Middle Anatolia  
(Ramme, 1951; Karabağ, 1958).

Remarks: This species was described by Ramme from  
Erciyes dağı in Kayseri province ca. 430 km. far from  
Seydişehir, the locality of my specimens. In the same  
paper, Ramme (1951: 165) described *Isophya*  
*anatolica* from Beyşehir that is only 34 km. far from  
Seydişehir. *I. anatolica* Ramme, 1951 has very similar  
cercus and subgenital plate with *I. pentheri* Ramme,  
1951. My specimens have exactly same cercus and  
subgenital plate in shape with *I. pentheri* Ramme,  
1951. But, for a better comparison more material is  
needed.

***Poecilimon zonatus* Bolivar, 1899**

*Poecilimon zonatus* Bolivar, 1899, Annls Soc. ent.  
Belg. 43: 597.

Material Studied: Konya: Bozkır, Karacaören,  
21.6.1997, 12 ♂, 4 ♀; 3 km. S. of Elmaağac, 1200m.  
13.6.1997, 11 ♂, 5 ♀, 2 ♀ nymphs (in alcohol).  
Distribution: Anatolia (Ramme, 1951; Bei-Bienko,  
1954; Karabağ, 1958).

***Poecilimon davisi* Karabağ, 1953**

*Poecilimon davisi* Karabağ, 1953, Eos, 29:183.

Material Studied: Konya: Bozkır, Kuruçay, 1400m.  
14.6.1997, 22 ♂, 14 ♀. Distribution: S. Anatolia  
(Karabağ, 1958).

***Parapoecilimon antalyensis* Karabağ, 1975**

*Parapoecilimon antalyensis* Karabağ, 1975, J. nat.  
Hist. London, 9:340.

Material Studied: Antalya: Alanya, Alarahan, 75m.  
21.6.1997, 3 ♂, 2 ♀; Kaplanhanı yaylası, 22.6.1997, 1  
♂, 1 ♀, 2 ♂ nymphs, 1 ♀ nymph. Distribution: S.  
Anatolia (Karabağ, 1975).

Remarks: After the original description of this  
interesting species, these are the first records from the  
localities, situated 40-90 km. east of the type locality.  
Known only from southern slopes of the Middle  
Taurus Mountains.

***Tettigoniinae******Decticus albifrons* (Fabricius, 1775)**

*Locusta albifrons* Fabricius, 1775, Syst.Ent., p.286.

Material Studied: Antalya: Alanya, Alarahan, 75m.  
21.6.1997, 2 ♀. Distribution: N. Africa, Europe,  
Anatolia, Caucasus, Middle East, S.W. Asia (Ramme,  
1951; Harz, 1969).

***Platycleis (s.str.) intermedia* intermedia (Serville,  
1839)**

*Decticus intermedium* Serville, 1839, Ins.Orth., p.488.

Material Studied: Konya: Bozkır, Yalnızca, 1100m.  
4.9.1997, 1 ♀; Koruanlı, 1450m. 4.9.1997, 1 ♀;  
Taşkent, 1540-1700m. 4.9.1997, 1 ♂. Distribution:  
Morocco, S. Europe, Anatolia, Middle East, Caucasus,  
Middle and W. Asia (Zeuner, 1941; Ramme, 1951;  
Harz, 1969)

***Platycleis (s.str.) affinis* affinis Fieber, 1853**

*Platycleis affinis* Fieber, 1853, Lotos, 3:150.

Material Studied: Konya: İçeriçumra, Yazibelen,  
5.9.1997, 1 ♂. Distribution: N. Africa, Europe,  
Anatolia, Middle East, Caucasus (Zeuner, 1941;  
Ramme, 1951; Harz, 1969).

***Platycleis (s.str.) grisea* grisea (Fabricius, 1781)**

*Locusta grisea* Fabricius, 1781, Spec.Ins., 1:359.

Material Studied: Konya: Bozkır, Pinarcık, 1100m.  
20.6.1997, 1 ♂; Avdan, 1300m. 20.6.1997, 1 ♂;  
Taşkent, Belpınaribeli, 1850m. 4.9.1997, 2 ♂.  
Distribution: S. Europe, Anatolia (Harz, 1969).

***Drymadusa limbata grandis* Karabağ, 1961**

*Drymadusa limbata grandis* Karabağ, 1961, Bull. Br.  
Mus. (Nat. Hist.) Ent. 11 (1): 11.

Material Studied: Antalya: Alanya, Alarahan, 75m.  
22.5.1997, 1 ♂; 10 km. N. of Mahmurlar, 22.6.1997, 1  
♂ nymph. Distribution: S. Anatolia and probably Syria  
(Karabağ, 1961).

***Parapholidoptera antaliae* Nadig, 1991**

*Parapholidoptera antaliae* Nadig, 1991, Bull. Soc.  
Ent. Suisse 64: 127.

Material Studied: Antalya: Alanya, Alarahan, 75m.,  
22.5.1997, 1 ♂, 1 ♀, 1 ♀ nymph. Distribution: S.  
Anatolia (Nadig, 1991).

***Sureyaella bella* Uvarov, 1934***Sureyaella bella* Uvarov, 1934, Eos, 10:54.

Material Studied: Konya: Kulu, 1000m. 19.6.1997, 3 ♂, 2 ♀ (nymphs). Distribution: Middle Anatolia (Uvarov, 1934; Karabağ, 1958).

**Gryllidae: Gryllinae*****Gryllus campestris* Linnaeus, 1758**

*Gryllus Acheta campestris* Linnaeus, 1758, Syst. Nat. (ed.10), 1:425.

Material Studied: Konya: Bozkır, Yalnızca, 1100m. 13.6.1997, 1 ♂. Distribution: N. Africa, Europe, Anatolia, Middle East (Ramme, 1951; Harz, 1969).

***Modicogryllus aff. frontalis* (Fieber, 1844)**

*Acheta frontalis* Fieber, 1844, Ent. Monogr. Prag, p.127.

Material Studied: Antalya: Alanya, Alarahan, 75m. 22.5.1997, 1 ♀. Distribution: Middle and E. Europe, Anatolia, Caucasus, W. Asia (Ramme, 1951; Harz, 1969).

Remarks: A single ♀ has been captured in the area. It is most similar with *M. frontalis* (Fieber). Of course it should be inspected with ♂ again.

**Pyrgomorphidae*****Pyrgomorpha guentheri* Burr, 1899**

*Pyrgomorpha grylloides* var. nov. *guentheri* Burr, 1899, J. linn. Soc. Lond. (zool.) 27:417

Material Studied: Konya: 6 km. N.E. of Bozkır, 1400m. 4.9.1997, 3 ♂, 1 ♀. Distribution: Anatolia, Caucasus, Middle East (Ramme, 1951; Bei-Bienko and Mistshenko, 1951; Karabağ, 1958).

**Pamphagidae: Akicerinae*****Glyphotmethis holtzi* aff. *pulchripes* (Uvarov, 1943)**

*Asiotmethis pulchripes* Uvarov, 1943, Trans. R. Ent. Soc. Lond. 98:57.

Material Studied: Konya: Bozkır, Avdan, 1300m. 20.6.1997, 3 ♂, 2 ♀; Pinarcık, 1100m. 20.6.1997, 4 ♂, 4 ♀; 6 km. N.E. of Bozkır, 1400m. 20.6.1997, 1 ♂, 5 ♀. Distribution: Middle Anatolia (Cejchan, 1965).

Remarks: Most species of the genus are found in Turkey only. Populations have great varieties. So, there are some identification problems among the species and the subspecies. The present specimens are more similar with this subspecies. Meanwhile, a revisional study on *Glyphotmethis* Bei-Bienko has been carrying out by me.

***Glyphotmethis holtzi spinosus* Karabağ, 1956**

*Glyphotmethis holtzi spinosus* Karabağ, 1956, Eos, 32:127.

Material Studied: Konya: Kulu, 1000m. 19.6.1997, 1 ♂, 1 ♀ nymph (leg. M. Ünal); S.W. shore of Tuz Gölü, 25.9.1996, 1 ♀ (leg. M. Vural). Distribution: Middle Anatolia (Cejchan, 1965).

**Pamphaginae*****Paranocarodes aff. fieberi* (Brunner von Wattenwyl, 1882)**

*Nocarodes fieberi* Brunner von Wattenwyl, 1882, Prodromus Eur. Orth., p.189.

Material Studied: Konya: Bozkır, Sorkun, 1100m. 14.6.1997, 1 ♂; Kuruçay, 1400m. 14.6.1997, 1 ♀; Antalya: Akseki, 1700m. 27.5.1997, 1 ♀. Distribution: Anatolia (Ramme, 1951; Bei-Bienko and Mistshenko, 1951; Demirsoy, 1973).

Remarks: Identification of the individuals of *Paranocarodes* Bolivar is rather difficult. 6 species and subspecies were described from Taurus Mountains. Of which, 3 species have been synonymized by Demirsoy (1973) in the last revision of the Turkish species. But the collected new material could not be determined by previous revisions (Bei-Bienko and Mistshenko, 1951; Ramme, 1951; Demirsoy, 1973). The genus should be reviewed that I have in mind.

**Acrididae: Catantopinae*****Sphenophyma rugulosa* (Stål, 1876)**

*Platynophyma rugulosa* Stål, 1876, K. svenska Vetensk Akad. Handl., 4 (5):18.

Material Studied: Antalya: Alanya, Mahmutlar, 22.6.1997, 1 ♀ nymph; 10 km. N. of Mahmutlar, 22.6.1997, 6 ♂, 5 ♀, 1 ♂ nymph, 5 ♀ nymphs. Distribution: Anatolia, Middle East (Ramme, 1951; Bei-Bienko and Mistshenko, 1951; Karabağ, 1958).

***Pezotettix anatolica* Uvarov, 1934**

*Pezotettix anatolica* Uvarov, 1934, Eos, 10:112.

Material Studied: Antalya: Alanya, Alarahan, 75m. 21.6.1997, 5 ♂, 7 ♀, 2 ♂ nymphs, 7 ♀ nymphs; Demirtaş, 50m. 22.6.1997, 3 ♂; 10 km. N. of Mahmutlar, 22.6.1997, 1 ♂. Distribution: Anatolia (Ramme, 1951; Bei-Bienko and Mistshenko, 1951; Karabağ, 1958).

**Eyprepocnemidinae*****Eyprepocnemis plorans* (Charpentier, 1825)**

*Gryllus plorans* Charpentier, 1825, Hor. Ent., p.134.

Material Studied: Antalya: Alanya, Alarahan, 75m. 21.6.1997, 3 ♂, 1 ♀ (nymphs).

Distribution: N. Africa, S. Europe, Anatolia, Middle East, W. Asia (Ramme, 1951; Bei-Bienko and Mistshenko, 1951; Harz, 1975).

### **Calliptaminae**

#### ***Calliptamus tenuicercis* Tarbinskyi, 1930**

*Calliptamus tenuicercis* Tarbinskyi, 1930, Bull. Akad. Sci. URSS, p.180, 183.

Material Studied: Konya: Bozkır, Korualan, 1450m. 4.9.1997, 1 ♀; 6 km. N.E. of Bozkır, 1400m. 5.9.1997, 1 ♂, 2 ♀; 10 km.W. of Hadim, 4.9.1997, 1 ♀; Hadim, İğdeören, 1300m. 4.9.1997, 1 ♂; Taşkent, 1540-1700m. 4.9.1997, 3 ♂, 3 ♀; Belpinaribeli, 1850m. 4.9.1997, 3 ♂; Dinek-Kavaklı, 1150m. 5.9.1997, 1 ♂, 3 ♀; İçeriçumra, Yazıbelen, 5.9.1997, 1 ♂, 1 ♀. Distribution: Anatolia, Middle East, Caucasus, S.W. Asia (Ramme, 1951; Bei-Bienko and Mistshenko, 1951; Jago, 1963; Harz, 1975).

#### ***Calliptamus barbarus* (Costa, 1836)**

*Acridium barbarum* Costa, 1836, Faun. Nap. Ortott., p.13.

Material Studied: Konya: İçeriçumra, Yazıbelen, 5.9.1997, 1 ♂, 4 ♀; 10 km. S. of İçeriçumra, 1000m. 1 ♂, 1 ♀; Antalya: Alanya, Alaran, 75m. 21.6.1997, 1 ♂, 2 ♀, 1 ♀ nymph.; Mahmurlar, 100m. 22.6.1997, 6 ♂, 1 ♀; Demirtaş, 50m. 22.6.1997, 1 ♂, 2 ♀. Distribution: Europe, Anatolia, Caucasus, Middle East, S.W. Asia (Ramme, 1951; Bei-Bienko and Mistshenko, 1951; Jago, 1963; Harz, 1975).

#### ***Calliptamus coelesyriensis* (Giglio-Tos, 1893)**

*Caloptenus coelesyriensis* Giglio-Tos, 1893, Bull. Mus. Torino, 8:10, 11.

Material Studied: Antalya: Alanya, 10 km. N. of Mahmurlar, 22.6.1997, 2 ♂, 1 ♀ nymph. Distribution: N.E. Africa, Anatolia, Middle East, Caucasus, Middle and W. Asia (Bei-Bienko and Mistshenko, 1951; Jago, 1963; Harz, 1975).

### **Aridinae**

#### ***Acrida bicolor* (Thunberg, 1815)**

*Truxalis bicolor* Thunberg, 1815, Mem.Acad.Petersb., 5:266.

Material Studied: Konya: İçeriçumra, 1000m. 3.9.1997, 1 ♂, 1 ♀; Dinek-Kavaklı, 1150m. 5.9.1997, 2 ♂, 1 ♀. Distribution: Africa, S. Europe, Anatolia, Middle East (Ramme, 1951; Bei-Bienko and Mistshenko, 1951; Dirsh, 1954; Harz, 1975).

### **Gomphocerinae**

#### ***Ramburiella turcomana* (Fischer de Waldheim, 1833)**

*Oedipoda turcomana* Fischer de Waldheim, 1833, Bull.Soc.Nat.Mosk., 6:351.

Material Studied: Antalya: Alanya, Mahmurlar, 100m. 22.6.1997, 2 ♀; 10km. N. of Mahmurlar, 22.6.1997, 2 ♀ nymphs. Distribution: S. Europe, Anatolia, Middle East, Caucasus, Middle and S.W. Asia (Ramme, 1951; Bei-Bienko and Mistshenko, 1951; Harz, 1975).

#### ***Arcyptera (Paracyptera) labiata* (Brulle, 1832)**

*Podisma labiata* Brulle, 1832, Exped.Sci.Moree, 3:95.

Material Studied: Konya: Bozkır, Karacaören, 21.6.1997, 2 ♂, 3 ♀; Kuruçay, 1400m. 14.6.1997, 7 ♂, 3 ♀; Hadim, 13.6.1997, 1200-1300m. 4 ♂, 2 ♀, 1 ♀ nymph. Distribution: S. Balkans and Anatolia (Harz, 1975; Willemse, 1984).

#### ***Dociostaurus (Kazakia) jagoi jagoi* Soltani, 1978**

*Dociostaurus (Kazakia) jagoi jagoi* Soltani, 1978, J.Ent.Soc.Iran, 2:30.

Material Studied: Antalya: Alanya, Demirtaş, 50m. 22.6.1997, 1 ♂, 6 ♀. Distribution: N. Africa, Anatolia, Middle East (Soltani, 1978).

#### ***Dociostaurus (Kazakia) brevicollis* (Eversmann, 1848)**

*Oedipoda brevicollis* Eversmann, 1848, Addit. Fischer de Waldheim, Orth.Ross., 11.

Material Studied: Konya: Bozkır, Kuruçay, 1400m. 14.6.1997, 1 ♂; Pinarcık, 1100m. 20.6.1997, 1 ♂. Distribution: Europe, Anatolia, Caucasus, Middle East, W. Asia, S. Siberia (Ramme, 1951; Bei-Bienko and Mistshenko, 1951; Harz, 1975; Soltani, 1978).

#### ***Dociostaurus (s.str.) salmani* Demirsoy, 1977**

*Dociostaurus (Dociostaurus) salmani* Demirsoy, 1977, Türk.Caeli.Faun., p.195.

Material Studied: Konya: Kulu, 1000m. 19.6.1997, 1 ♂, 4 ♀. Distribution: Anatolia (Demirsoy, 1977; Çiplak, 1991; Ünal, 1995).

#### ***Dociostaurus (Stauronotulus) cappadocicus* (Azam, 1913)**

*Stauronotus crassiusculus* var. *cappadocicus* Azam, 1913, Bull. Soc. ent.Fr. p.219.

Material Studied: Konya: Bozkır, Karacaören, 21.6.1997, 3 ♂; 6 km. N.E. Of Bozkır, 1400m. 15.6.1997, 1 ♂, 1 ♀ nymph; 20.6.1997, 2 ♂, 1 ♀; 3km. S. of Elmaağaç, 1200m. 13.6.1997, 6 ♂, 5 ♀ (in alcohol). Distribution: Anatolia (Ramme, 1951; Demirsoy, 1977).

***Dociostaurus (Stauronotulus) hauensteini (Bolivar, 1893)***

*Stauronotus hauensteini* Bolivar, 1893,  
Rev.Bio.Nord.France, 5:480.

Material Studied: Konya: Kulu, 1000m., 19.6.1997, 2 ♂, 4 ♀. Distribution: Anatolia, Middle East, W. Asia (Ramme, 1951; Soltani, 1978).

***Dociostaurus (Notostaurus) anatolicus (Krauss, 1896)***

*Stauronotus anatolicus* Krauss, 1896, Zol.Jahrb.Syst., 9:560.

Material Studied: Konya: İçeriçumra, Yazibelen, 5.9.1997, 2 ♂, 3 ♀. Distribution: Balkans, Anatolia, Middle East, Caucasus, W. Asia (Ramme, 1951; Bei-Bienko and Mistshenko, 1951; Harz, 1975; Soltani, 1978).

***Stenobothrus (s.str.) fischeri (Eversmann, 1848)***

*Oedipoda fischeri* Eversmann 1848, Addit. Fischer de Waldheim, Orth.Ross., 11.

Material Studied: Konya: Bozkır, Kuruçay, 1400m. 14.6.1997, 1 ♂, 2 ♀; Antalya: Akseki, 21.6.1997, 3 ♂, 4 ♀. Distribution: Europe, Anatolia, Caucasus, Middle and W. Asia (Ramme, 1951; Bei-Bienko and Mistshenko, 1951; Harz, 1975).

***Stenobothrus (s.str.) zubowskyi Bolívar, 1899***

*Stenobothrus zubowskyi* Bolívar, 1899,  
Ann.Soc.Ent.Belg., 43:588.

Material Studied: Konya: 6 km. N.E. of Bozkır, 15.6.1997, 1 ♂, 1 ♀; 20.6.1997, 1 ♂, 5 ♀; Pınarcık, 1100m. 20.6.1997, 4 ♂, 3 ♀; 15km. E of Aydinkelşla, 15.6.1997, 2 ♂; Kuruçay, 1400m. 14.6.1997, 1 ♂; Avdan, 1300m. 20.6.1997, 1 ♂; Sorkun, 1200m. 14.6.1997, 1 ♂, 3 ♀ (in alcohol); Hadim, 1200-1300m. 13.6.1997, 1 ♂. Distribution: Anatolia, Caucasus (Ramme, 1951; Bei-Bienko and Mistshenko, 1951).

Remarks: These are southermost populations in Turkey.

***Omocestus petraeus (Brisout, 1855)***

*Acridium petraeum* Brisout, 1855, Bull. Soc. ent. Fr. 3 (3): 114.

Material Studied: Konya: 6 km. N.E. of Bozkır, 1400m. 5.9.1997, 2 ♀; 10 km. W. of Hadim, 4.9.1997, 1 ♂, 2 ♀; Taşkent, 1540-1700m. 4.9.1997, 1 ♂, 4 ♀. Distribution: Europe, Anatolia, Caucasus, S. Siberia (Ramme, 1951; Bei-Bienko and Mistshenko, 1951; Harz, 1975).

***Chorthippus (s.str.) parallelus parallelus (Zetterstedt, 1821)***

*Gryllus parallelus* Zetterstedt, 1821, Orth.Suec., p.85.

Material Studied: Konya: 3 km. S. of Seydişehir, 1000m. 14.6.1997, 2 ♂. Distribution: Europe, Anatolia, Caucasus, N.W. Asia, Siberia (Ramme, 1951; Bei-Bienko and Mistshenko, 1951; Harz, 1975).

***Chorthippus (s.str.) dichrous (Eversmann, 1859)***

*Oedipoda dichroa* Eversmann, 1859, Bull. Soc. Nat. Mosc. 31(1):132.

Material Studied: Konya: Bozkır, Dere, 1100m. 20.6.1997, 4 ♂, 3 ♀; Akçapınar, 1200m. 3.9.1997 1 ♂, 1 ♀; 10km. W. of Hadim, 4.9.1997, 1 ♂, 1 ♀; Taşkent, Belpinaribeli, 1850m. 4.9.1997, 5 ♂, 2 ♀; İçeriçumra, 1000m. 3.9.1997, 10 ♂, 1 ♀. Distribution: Middle Europe, Balkans, Anatolia, Caucasus, Middle and W. Asia, W. Siberia (Bei-Bienko and Mistshenko, 1951; Harz, 1975).

***Chorthippus (Glyptobothrus) aff. brunneus (Thunberg, 1815)***

*Gryllus brunneus* Thunberg, 1815, Mem. Acad. Petersb., 5:249.

Material Studied: Konya: 6 km. N.E. of Bozkır, 1400m. 20.6.1997, 2 ♂, 1 ♀; Dere, 1100m. 20.6.1997, 1 ♂, 2 ♀; Avdan, 1300m. 20.6.1997, 1 ♂, 1 ♀; Sorkun, 1200m. 14.6.1997, 2 ♂ (in alcohol); 3 km. S. of Elmağaç, 1200m. 13.6.1997, 2 ♂, 2 ♀ (in alcohol); Hadim, 1200-1300m. 13.9.1997, 2 ♂; Antalya: Alanya, 10km. N. of Mahmutlar, 22.6.1997, 2 ♂, 2 ♀. Distribution: Europe (Ragge et al., 1990).

Remarks: This species was known from N. Africa and W. Europe to E. Asia and Siberia (Bei-Bienko and Mistshenko, 1951; Harz, 1975). But in the result of acoustics studies (Ragge et al., 1990), it does not occur in Africa. Known from Europe except Iberian and Balkan peninsula. Thus, the occurrence of this species in Turkey is doubtful. But it must be reviewed using their songs in the localities that were reported in Turkey.

***Chorthippus (Glyptobothrus) biguttulus (Linnaeus, 1758)***

*Gryllus Locusta biguttulus* Linnaeus, 1758, Syst. Nat. (ed.10), 1:433.

Material Studied: Konya: 6 km. N.E. of Bozkır, 1400m. 5.9.1997, 2 ♀; Dere, 1100m. 4.9.1997, 2 ♂, 1 ♀; Pınarcık, 1100m. 20.6.1997, 1 ♀; Avdan, 1300m. 20.6.1997, 1 ♂; Karacaören, 21.6.1997, 1 ♀; Çağlayan, 1000m. 14.6.1997, 1 ♂, 2 ♀; Sorkun, 1200m. 14.6.1997, 1 ♀ (in alcohol); Yalnızca, 1100m. 13.6.1997, 1 ♀; Dinek-Kavaklı, 1150m. 5.9.1997, 1 ♀; 10km. W. of Hadim, 4.9.1997, 1 ♂, 3 ♀; Taşkent, 1540-1700m. 4.9.1997, 2 ♂; 3 km. S. of Seydişehir, 1000m. 14.6.1997, 1 ♂. Distribution: It was known from N. Africa, Europe, Anatolia, Caucasus, N. Asia, Siberia (Bei-Bienko and Mistshenko, 1951; Harz, 1975). But now, known from Middle and N. Europe (Ragge et

al., 1990). A revision is needed with the analyses of the songs in the other older localities.

Remarks: Only nominate subspecies was reported from Turkey so far. Helversen (1989) has found a new subspecies from S. Balkans and N.W. Anatolia, *C. biguttulus euhedickei* Helversen, in the result of analysis the calling songs. Also the other populations need analyses of the songs in Turkey. Probably new taxa can be discovered by using this method.

#### ***Dasyhippus escalerae (Bolívar, 1899)***

*Gomphecerus escalerae* Bolívar, 1899, Annls Soc. ent. Belg. 43:590.

Material Studied: Konya: Bozkır, Yalnızca, 1100m. 13.6.1997, 1 ♂, 2 ♀. Distribution: Anatolia (Ramme, 1951; Nasracki & Ünal, 1995).

#### ***Oedipodinae***

##### ***Aiolopus thalassinus (Fabricius, 1781)***

*Gryllus thalassinus* Fabricius, 1781, Spec. Ins., 1:367.

Material Studied: Antalya: Alanya, Alarahan, 75m., 21.6.1997, 1 ♀. Distribution: Africa, Europe, Anatolia, Middle East, Caucasus, Middle and S.W. Asia (Ramme, 1951; Bei-Bienko and Mistshenko, 1951; Hollis, 1968; Harz, 1975).

##### ***Aiolopus strepens (Latreille, 1804)***

*Acridium strepens* Latreille, 1804, Hist. Nat. Crust. Ins., 12:154.

Material Studied: Antalya: Alanya, Alarahan, 75m., 21.6.1997, 1 ♀. Distribution: N. Africa, Middle and S. Europe, Anatolia, Caucasus, Middle East, Middle and W. Asia (Ramme, 1951; Bei-Bienko and Mistshenko, 1951; Hollis, 1968; Harz, 1975).

##### ***Locusta migratoria (Linnaeus, 1758)***

*Gryllus Locusta migratorius* Linnaeus, 1758, Syst. Nat. (ed.10), 1:432.

Material Studied: Konya: Taşkent, Belpınarıbeli, 1850m. 4.9.1997, 1 ♂ nymph, 1 ♀. Distribution: Anatolia, Caucasus, Middle and W. Asia (Ramme, 1951; Bei-Bienko and Mistshenko, 1951; Harz, 1975).

##### ***Pyrgodera armata (Fischer de Waldheim, 1820)***

*Acridium armatum* Fischer de Waldheim, 1820, Ent. Ross., 1:37.

Material Studied: Konya: Bozkır, Dere, 1100m. 20.6.1997, 3 ♀; Karacaören, 21.6.1997, 5 ♂, 2 ♀; Çağlayan, 1000m. 14.6.1997, 1 ♂; Sorkun, 1200m. 14.6.1997, 1 ♂, 1 ♀ (in alcohol); 3 km. S. of Elmaağaç, 1200m. 13.6.1997, 1 ♂ (in alcohol); Antalya: Alanya, Çayarası, 22.6.1997, 1 ♀. Distribution: Anatolia, Caucasus, Middle East, Middle and W. Asia (Ramme, 1951; Bei-Bienko and Mistshenko, 1951; Harz, 1975).

#### ***Oedipoda aurea Uvarov, 1923***

*Oedipoda miniata* ssp. *aurea* Uvarov, 1923, Entomologist mon. Mag. 11 (3): 32.

Material Studied: Konya: 6 km. N.E. of Bozkır, 1400m. 5.7.1997, 2 ♀. Distribution: Balkans, Anatolia, Caucasus, Middle East (Ramme, 1951; Bei-Bienko and Mistshenko, 1951; Harz, 1975).

#### ***Oedipoda miniata miniata (Pallas, 1771)***

*Gryllus miniatus* Pallas, 1771, Reis. Russ. Reich., 1:467.

Material Studied: Konya: Bozkır, Dere, 1100m. 20.6.1997, 1 ♂, 1 ♀; Yalnızca, 1100m. 4.9.1997, 1 ♀, 1 ♀ nymph; 5 km E. of Aydınışla, 1400m. 5.9.1997, 5.9.1997, 2 ♂; Yazibelen, 5.9.1997, 1 ♀; Hadim, Korualan, 1450m. 4.9.1997, 2 ♂; İğdeören, 1300m. 4.9.1997, 3 ♂, 1 ♀; Dinek-Kavaklı, 1150m. 5.9.1997, 2 ♂, 1 ♀; Antalya: Alanya, Mahmutlar, 100m. 22.6.1997, 2 ♀; Demirtaş, 50m. 22.6.1997, 2 ♂, 4 ♀. Distribution: N. Africa, E. and S. Europe, Anatolia, Caucasus, Middle East, Middle and W. Asia, S.W. Siberia (Ramme, 1951; Bei-Bienko and Mistshenko, 1951; Harz, 1975).

#### ***Oedipoda caerulescens (Linnaeus, 1758)***

*Gryllus Locusta caerulescens* Linnaeus, 1758, Syst. Nat. (ed.10), 1:432.

Material Studied: Konya: Bozkır, Dere, 1100m. 4.9.1997, 1 ♂; 10km. W. of Hadim, 4.9.1997, 1 ♀; Taşkent, 1540-1700m. 4.9.1997, 2 ♂, 3 ♀. Distribution: N. Africa, Europe, Anatolia, Caucasus, Middle East, Middle and S.W. Asia (Ramme, 1951; Bei-Bienko and Mistshenko, 1951; Harz, 1975).

#### ***Acrotylus insubricus insubricus (Scopoli, 1786)***

*Gryllus insubricus* Scopoli, 1786, Del. Fau. Flor. Insubr., 1:64.

Material Studied: Konya: Taşkent, 1540-1700m. 4.9.1997, 1 ♂; İçeriçumra, 1000m. 3.9.1997, 1 ♂; 10 km. S. of İçeriçumra, 1000m. 5.9.1997, 1 ♂, 1 ♀; Yazibelen, 5.9.1997, 1 ♂; Antalya: Akseki, 1700m. 24.5.1997, 1 ♂, 1 ♀. Distribution: E. Europe, Balkans, Anatolia, Caucasus, Middle East, S.W. Asia (Ramme, 1951; Bei-Bienko and Mistshenko, 1951; Harz, 1975).

#### ***Pseudocleles ledereri (Saussure, 1884)***

*Thalpomena ledereri* Saussure, 1884, Mem. Soc. Geneve 28:184.

Material Studied: Konya: Taşkent, Belpınarıbeli, 1850m. 4.9.1997, 17 ♂, 9 ♀. Distribution: Anatolia (Ramme, 1951; Bei-Bienko and Mistshenko, 1951; Karabağ, 1958; Demirsoy, 1977).

Remarks: This interesting species has been found from the upper heights of the Taurus.

***Sphingonotus (s.str.) theodori theodori* Uvarov, 1923**  
*Sphingonotus coeruleans theodori* Uvarov, 1923, Bull. Soc. R. Ent. Egypte, p.195.

Material Studied: Konya: Bozkır, Boyalı, 1300m. 5.9.1997, 1 ♀; Yalnızca, 1100m. 4.9.1997, 1 ♀; 5 km. E. of Aydinkelışla, 1400m. 5.9.1997, 3 ♂; 10km. S. of İçeriçumra, 1000m. 5.9.1997, 2 ♂; Yazibelen, 5.9.1997, 3 ♂, 3 ♀. Distribution: Anatolia, Middle East, Sinai (Mistshenko, 1936; Ramme, 1951; Bei-Bienko and Mistshenko, 1951).

***Sphingonotus (s.str) pilosus* Saussure, 1884**

*Sphingonotus pilosus* Saussure, 1884, Mém. Soc. Genéve, 28(9):201.

Material Studied: Konya: 6 km. N.E. of Bozkır, 1400m. 5.9.1997, 1 ♂, 3 ♀; 5 km. E. of Aydinkelışla, 1400m. 5.9.1997, 2 ♂, 1 ♀. Distribution: Anatolia, Caucasus, S.W. Asia (Mistshenko, 1936; Ramme, 1951; Bei-Bienko and Mistshenko, 1951).

***Sphingonotus (s.str.) nebulosus anatolicus* Uvarov, 1930**

*Sphingonotus nebulosus anatolicus* Uvarov, 1930, Eos, 6:366.

Material Studied: Konya: İçeriçumra, Yazibelen, 5.9.1997, 1 ♂, 2 ♀; Taşkent, 1540-1700m. 4.9.1997, 4 ♂, 5 ♀. Distribution: Anatolia (Mistshenko, 1936; Ramme, 1951; Bei-Bienko and Mistshenko, 1951).

Remarks: The basal part of hind wings are colourless in the population of Yazibelen.

***Sphingonotus (s.str.) turcicus turcicus* Uvarov, 1930**

*Sphingonotus turcicus* Uvarov, 1930, Eos, 6:367.

Material Studied: Konya: Bozkır, Boyalı, 1300m. 5.9.1997, 2 ♀; Taşkent, 1540-1700m. 4.9.1997, 4 ♂, 2 ♀; Dineksaray, 1100m. 3.9.1997, 1 ♀; 10km. S. of İçeriçumra, 1000m. 5.9.1997, 2 ♂; Yazibelen, 5.9.1997, 1 ♀. Distribution: Anatolia (Mistshenko, 1936; Ramme, 1951; Bei-Bienko and Mistshenko, 1951).

Remarks: Hind wings with colourless basal part in a single ♀ from Dineksaray.

## References

- Bei-Bienko, G.Ya and Mistshenko, L.L., 1951, *Fauna of the U.S.S.R. Locusts and Grasshoppers of the U.S.S.R. and Adjacent Countries*. Part 1. Zoological Institute of the U.S.S.R. Academy of Sciences, 38: 400.
- Bei-Bienko, G.Ya and Mistshenko, L.L., 1951, *Fauna of the U.S.S.R. Locust and Grasshoppers of the U.S.S.R. and Adjacent Countries*. Part 2. Zoological Institute of the U.S.S.R. Academy of Sciences, 40: 291.
- Bei-Bienko, G.Ya., 1954, *Fauna of the U.S.S.R. Phaneropterinae (Tettigonoidea: Orthoptera)*. Zoological Institute of the U.S.S.R. Academy of the Sciences., 59:375.
- Cejchan, A., 1965, A Preliminary revision of the genus *Glyphotmethis* Bey-Bienko (Orthoptera: Pamphagidae). *Ent. Mus. nat. Pragae*, 36:451-490.
- Demirsoy, A., 1973, Revision der anatolischen Pamphaginae (Saltatoria, Caelifera, Pamphagidae). *Ent. Mitt. zool. Mus. Hamburg*, 4(3):403-428.
- Demirsoy, A., 1977, *Türkiye Caelifera (Insecta,Orthoptera) faunasının tesbiti ve taksonomik incelenmesi* (1). Atatürk Üniversitesi Fen Fak. yayınları, 80:1-25.
- Gümüşsuyu, İ., 1981, *Türkiye Gryllidae (Orthoptera) faunası üzerine sistematik çalışmalar ile türlerin habitat ve davranışlarına ait gözlemler*. Ankara Bölge Zirai Mücadele Araştırma Enstitüsü, araştırma eserleri serisi, no: 42.
- Harz, K., 1969, *The Orthoptera of Europe (Die Orthopteren Europas)*. Hague, 1:1-749.
- Harz, K., 1975, *The Orthoptera of Europe (Die Orthopteren Europas)*. Hague, 2:1-939.
- Helversen, v.O., 1989, Bemerkungen zu *Chorthippus biguttulus hedickei* (Ramme, 1942) und Beschreibung von *Chorthippus biguttulus eu hedickei* n.ssp. *Articulata*, 4:26-34.
- Karabağ, T., 1950, Five new species of *Poecilimon* Fischer (Orthoptera, Tettigoniidae) from Turkey. *Proc. R. ent. London*, 19: 157-161.
- Karabağ, T., 1956, Some new and less known Acrididae from Turkey. *Eos*, 32:125-134.
- Karabağ, T., 1958, *Türkiye'nin Orthoptera faunası (The Orthoptera Fauna of Turkey)*. Ankara Univ. Fen Fak. yay. 81: 1-198.
- Karabağ, T., 1964, Some new species and new records of Tettigoniidae (Orthoptera) from Turkey. *Communications*, 13: 36-55.
- Karabağ, T., Gümüşsuyu, İ., Balamir, S. ve Tutkun, E., 1971, Türkiye Orthoptera Faunası'nın Tesbiti Üzerine Araştırmalar. *Bitki Koruma Bülteni*, 11(2): 73-98.
- Karabağ, T., Gümüşsuyu, İ., Balamir, S. ve Tutkun, E., 1974, Türkiye Orthoptera Faunası'nın Tesbiti Üzerine Araştırmalar. *Bitki Koruma Bülteni*, 14 (1): 2-18.
- Karabağ, T., 1975, Studies in the Turkish Orthoptera (Insecta). I. New Species and less known Tettigoniidae. *J. nat. Hist.*, 9: 337-350.
- Karabağ, T., Gümüşsuyu, İ. ve Tutkun, E., 1980, Türkiye Orthoptera faunasının tesbiti üzerine araştırmalar (III). *Bitki Koruma Bülteni*, 20 (1-4): 1-25.
- Mistshenko, L., 1936. XII. Revision of Palaearctic species of the genus *Sphingonothus* Fieber (Orth. Acrid.). *Eos*, 12:65-282.
- Nadig, A., 1991, Eine neue Parapholidoptera-Art aus der südlichen Türkei: *Parapholidoptera antaliae* spec. nova (Orthoptera, Tettigoniidae). *Mitt. schweiz. Ent. Ges.* 64: 127-131.
- Naskrecki, P. & Ünal, M., 1995, The Orthoptera of Hatay province, S.Turkey. *Beitr. Ent.* 45 (2): 393-419.
- Ragge, D.R., W.J. Reynolds and F. Willems, 1990, The songs of the European grasshoppers of the Chorthippus biguttulus group in relation to their taxonomy, speciation and biogeography (Orthoptera: Acrididae). *Bol. San. Veg. Plagas (Fuera de serie)*, 20:239-245.

- Ramme, W., 1951. Zur Systematik Faunistik und Biologie der Orthopteren von Südost Europa und Vorderasien. *Mitt. zool. Mus. Berlin*, 27:1-431.
- Uvarov, B.P., 1934. Studies in the Orthoptera of Turkey, Iraq and Syria. *Eos*, 10:21-119.
- Ünal, M., 1997. Nevşehir ili Avanos çevresinin Orthopter'lerinin Eko-faunası üzerine araştırmalar. *Priamus*, 9 (1): 1-54.
- Ünal, M., 1999. Notes on Orthoptera of Western Turkey, with Description of a new genus and four new Species. *J. Orthoptera. Res.*, 8: 243-255.
- Weidner, H., 1969. Beiträge zur Kenntnis der Feldheuschrecken (Caelifera) Anatoliens. *Mitt. Zool. Mus. Hamburg.*, 66:145-226.
- Werner, F., 1901. Die Dermapteren und Orthopterenfauna Kleinasiens. *Sitzb. Akad. Wiss. Mathem.-nat.*, 110: 259-306.
- Willemse, F., 1973. A study of the genus Paracaloptenus I. Bolivar, 1876 (Orthoptera, Acridoidea, Calliptaminae). *Beaufortia* 20 (270): 179-192.
- Zeuner, F.E., 1941. The classification of the Decticinae hitherto included in Platycleis Fieb. or Metrioptera Wesm. (Orthoptera, Saltatoria). *Trans. R. ent. Soc. London* 91: 1-49.

\*\*\*\*\*

**Contents / İçindekiler:** Ünal, M. Contributions to the *Orthoptera fauna* of the Middle Taurus Mountains (South Turkey) s.1 - Editorial, s.8.

**MISCELLANEOUS PAPERS****ISSN 1015-8235**

Miscellaneous Papers is the second international serial of the Centre for Entomological Studies Ankara (Cesa). It includes original papers of the research workers of the Centre on ecology, fauna, flora, distribution, check-list, taxonomy, nomenclature and morphology, as well as information and announcements from the Centre. The publication language are Turkish, Uighurian, English, German and Chinese.

**CENTRE FOR ENTOMOLOGICAL STUDIES ANKARA**

( A scientific Consortium)

(co-operation of a number of research workers for pure-scientific, not commercial purpose)

Web Page of the Cesa: [http://www.members.tripod.com/Cesa\\_1988/on.html](http://www.members.tripod.com/Cesa_1988/on.html)

Scientific Serials: Priamus (ISSN 1015-8243), Miscellaneous Papers (ISSN 1015-8235), Memoirs (ISSN-8227)

News of the month, an electronical magazine of the Cesa in the internet.

<http://www.members.tripod.com/entlep/News5.htm>

Owner / Sahibi - Editor / Yayıncı: Prof. Dr. Ahmet Ömer Koçak - Editor Assistents: Dr. Selma Seven, Dr. Muhabbet Kemal, M.Sc. Emine Demir.

Editorial Board of all Scientific Serials / Bütün Bilimsel Yayınların Yayın Kurulu: Insecta, taxonomy, nomenclature, ecology, faunistics: Prof. Dr. Ahmet Ömer Koçak (Turkey), Dr. Selma Seven (Turkey), Dr. Muhabbet Kemal, Yrd. Doç. Dr. Yusuf Hüseyinoğlu (Turkey), Mr. Günter Ebert (Germany), Dr. Thomas Meineke (Germany). Homoptera: M.Sc. Emine Demir (Turkey). Orthoptera: Dr. Piotr Naskrecki (U.S.A.), Dr. Mustafa Ünal (Turkey). Coleoptera / Chrysomelidae: Ass. Prof. M.S.Mohammedsaid (Malaysia). - Plant taxonomy, flora and vegetation: Prof. Dr. Mecit Vural (Turkey), Doç. Dr. Nezaket Adıgüzel (Turkey).

ALL RIGHTS RESERVED

Correspondences should be addressed to: Prof. Dr. Ahmet Ömer Koçak, 100. Yıl University, Fen-Edebiyat Fakültesi Biyoloji Bölümü, Kampus, Van / Turkey. e-mail: [a\\_kocak@mailcity.com](mailto:a_kocak@mailcity.com)

MISCELLANEOUS PAPERS AYDA BİR  
YAYINLANIR

Yüzüncü Yıl Üniversitesi Matbaasında bastırılmıştır.

All serials are recorded regularly by the Zoological Record, Biosis, Garforth House, 54 Micklegate, York, North Yorkshire. fax (01904) 612793 - [DCS@york.biosis.org](mailto:DCS@york.biosis.org)